

## STATURE ESTIMATION BY USING PER-CUTANEOUS TIBIAL LENGTH

<sup>1</sup>Mahima Srivastava, <sup>2</sup>Abhijeet Soni, <sup>3</sup>Anupam Johry

<sup>1</sup>Sr.Professor, Department of Anatomy, <sup>2</sup>P.G. Resident, Department of Forensic Medicine,

<sup>3</sup>Assistant Professor, Department of Forensic Medicine J.L.N. Medical College, Ajmer, Rjasthan

### ABSTRACT

**BACKGROUND:** Stature estimation is an important part to establish the individuality of an unidentified dead, body or any mutilated part of such body by the Medico-legal expert.

**AIM:** The present study made an attempt to establish the statistical correlation between stature and per-cutaneous tibial length (PCTL) by formulating a simple regression equation and multiplication factor (M.F.) for the people of Ajmer region.

**METHOD:** A random sample of 250 male and 250 female subjects of the Ajmer Region between the age group of 18-21 years were chosen. PCTL of right and left side were measured with the help of spreading caliper. Stature was estimated from PCTL statistically using simple regression analysis and M.F.

**RESULT:** On computing the data, the mean PCTL for male was found to be  $38.24 \pm 2.343$ cm, which was significantly ( $p < 0.0001$ ) greater than the female ( $36.064 \pm 2.464$ cm). The observed height was  $164.5 \pm 8.257$ cm and  $155.3 \pm 5.854$ cm for the male and female respectively. The regression formula derived for the male was  $y_0 = 105.971 + 1.53 \times (\text{PCTL}) \pm 7.452$  and for the female was  $y_0 = 103.76 + 1.43 \times (\text{PCTL}) \pm 4.69$ . The M.F was 4.302 for male and 4.306 for female. A significant positive correlation exists between the stature and PCTL using simple regression analysis and M.F.

**CONCLUSION:** It was concluded that the stature in Ajmer region can be determined by using the formula derived from the present study fairly accurately.

**KEY-WORDS:** Forensic Anthropology, Stature, Percutaneous Tibial Length, Height Estimation, Simple Regression Equation.

•-----•  
**Corresponding Authour:** Dr. Abhijeet Soni, P.G. Resident, J.L.N. Medical College, Ajmer  
Email-annu\_johry@yahoo.co.in  
•-----•

## INTRODUCTION

Stature estimation is an important part of the identification in human skeletal remains or body parts.<sup>1-4</sup> The estimation of stature from femur and tibia, are more accurate than the humerus and ulna,<sup>5</sup> especially the tibia and the femur as these have a direct correlation to the height of an individual.<sup>6</sup> Forensic While dealing with skeletal remains use of anatomical method for stature reconstruction has limited role due to non-availability of the complete skeleton from a scene of crime in most of the cases.<sup>7-8</sup> The mathematical method can be used in these cases where only a part of the body<sup>9</sup> or part of the bone<sup>10-12</sup> are available for determination of the stature. The lower limb length is the greatest contributor to the standing height, hence the most predictive equation is based on length of lower limb, the femur, Tibia and fibula.<sup>13-16</sup> The tibia is ideal in this application as it resists erosion and keeps its anatomical shape for long even after burial.<sup>17</sup> Tibia accounts for 22% of the total body length.<sup>18</sup>

Bone and stature of an individual are influenced by numerous factors as age, gender, race, geographical climate, nutritional and genetic factors.<sup>19-23</sup> Hence, the correlation factors of one region will not hold good for the other and separate research is required for that particular region.<sup>16,22-24</sup> There are various ways to estimate stature from bones, but the most easiest and reliable method is by regression analysis.<sup>22,25-26</sup> Regression formulae derived from the major long bones are generally considered to be more accurate. However the formulae derived cannot be generalized to all population groups, hence it is necessary to derive

regression equations which are region wise and population specific.<sup>23</sup>

## MATERIAL AND METHOD

**Study Design:** the present study is a cross Sectional study.

**Selection criteria:** A random sample of 500 subjects including 250 Male and 250 female, above the age of 21 years of the Ajmer region were selected for the present study, as bony growth is completed above 21yrs of age.<sup>27</sup> The following parameters were noted- Age, Gender, Height in cms (crown heel length), Percutaneous Tibial Length (PCTL) of right and left side in cms.

**Exclusion criteria:** Subjects who had a history of major trauma or fracture of the leg, Achondroplasia or any other congenital or hereditary bony disease was excluded from the study.

**Methodology:** All the measurements were taken by the same observer and with the same instrument, to avoid any technical and/or inter-observer error and to maintain reproducibility. The measurements were taken three times and their mean value was considered for estimation of height. Standing Height (Stature) of the subject was measured in a standing position on a standard Stadiometer with both feet in close contact with each other with the trunk straight along the vertical board, and the head adjusted in Frankfurt plane. All the measurement was taken in centimeters. For measuring the tibial length (PCTL) subject was asked to stand and keep his/her foot on a stool to maintain the angle between the flexor surface of leg and that of the thigh at

90°. Then two points were marked by skin marking pencil. Upper point - The medial most point on the upper border of medial condyle of the tibia and Lower point of the Tip of medial malleolus of the tibia. Distance between two points was measured with the help of Spreading Caliper to determine tibial length. The data were computed, tabulated and statistically analyzed using SSP 2015 Graph Pad Prism and Microsoft Excel Windows 2007 softwares. The data obtained were compared with the other similar studies.

## RESULTS

The statistical analysis of PCTL of the right and left side of the tibia in male and female was shown in Table 1. There was no significant difference ( $p > 0.05$ ) in the per-cutaneous length of the right and left tibia in both genders, thus showing bilateral symmetry in the length of Tibia in both genders. The mean PCTL for male was 38.24cm and for female was 36.06cm.

**Table 1:** Descriptive statistics of right and left side of Tibial length

Statistics Tibia	Male (n = 250)		Female (n = 250)	
	Rt PCTL	Lt PCTL	Rt PCTL	Lt PCTL
Range	34-45.8 cm	34-43.7 cm	32-42cm	32-48 cm
Mean	38.26cm	38.22cm	36.10cm	36.03cm
Std. Deviation	2.451	2.293	2.429	2.617
Std. Error	0.1492	0.1396	0.1479	0.1592
Coefficient of variation (CV)	6.41%	6.00%	6.73%	7.26%
t - value	t=0.1795 df=538		t=0.3170 df=538	
p - value	0.8576		0.7513	
P value summary	Ns		Ns	
Average mean (rt+lt)	38.24 cm		36.06 cm	

PCTL= per-cutaneous tibial length; Rt= right; Lt= left; Ns= not significant; df= degree of freedom.

The present study revealed that the standing height of many individuals were same, but their PCTL differed, i.e. the contribution of tibial length to the stature of a person varied from person to person, even for a given height.<sup>23</sup> Keeping this in view, Mean of stature and PCTL were taken into consideration and the data were calculated and analyzed (Table 2). The observed mean height was 164.5cm and 155.3cm; and mean PCTL was 38.24cm and 36.06cm in male and female respectively, which was significantly ( $p < 0.0001$ ) greater for male compared with female.

**Table 2:** Descriptive statistics of observed Height and Tibial length of male and female

Statistics	Male		Female	
	Height	PCTL	Height	PCTL
Range	143-182.5cm	34 - 43.65cm	147-176.5cm	32 - 44cm
Mean	164.5cm	38.24cm	155.3cm	36.06cm
Std. Deviation	8.257	2.343	5.854	2.464
Std. Error	0.5025	0.1426	0.3562	0.1499
Coefficient of variation (CV)	5.02%	6.13%	3.77%	6.83%
Student t- test between male and female Tibial length:				
t - value	t=10.50			
p - value	P<0.0001			
P value summary	***Significant			
Difference between means	2.176 ± 0.2069			
Are means signif. different? (P < 0.05)	Yes			
95% confidence interval	1.767 to 2.578			
R squared	0.1701			
Average mean PCTL(M+F)	37.151cm			

In Table 3, Correlation coefficients (r) of height and PCTL for male and female were 0.4342 and 0.6014 respectively, which were statistically significant. Since

there was a high correlation between the height and PCTL, a simple regression analysis was done between them for males and females and a simple regression formula was derived to predict the height from PCTL. The regression formula derived for male was  $y_0=105.971+1.53 \times (\text{PCTL}) \pm 7.452$

and for female was  $y_0= 103.76+1.43 \times (\text{PCTL}) \pm 4.69$ . The predicted height (y) derived from this formula was acceptable within a range of error and was in close approximation with that of the observed height.

**Table 3:** Formulation of Regression equation for calculating the stature from PCTL in male and female

Regression Statistics of Tibia	Male (observed ht=164.5cm)			Female (observed ht=155.3cm)		
	Rt	Lt	Average PCTL(rt+lt)	Rt	Lt	Average PCTL(rt+lt)
Independent variable(x) = PCTL	x1 = 38.26	x2 = 38.22	x0 = 38.24	x1 = 36.10	x2 = 36.03	x0 = 36.064
Intercept (a)	110.76	103.712	105.971	105.724	106.64	103.76
Regression coefficient (b)	1.404	1.59	1.53	1.373	1.35	1.43
Correlation coefficient(r)	0.4168	0.4416	0.4342	0.5699	0.6036	0.6014
Coefficient of determination (R <sup>2</sup> )	0.174	0.195	0.189	0.325	0.364	0.362
Std. error of estimate (SEE)	7.52	7.422	7.452	4.82	4.68	4.69
Significance (p)	***	***	***	***	***	***
Regression formula (y = a+bx)	y1= 110.76+ 1.404 (x)x1	y2= 103.712 + 1.59 (x) x2	y0=105.971+ 1.53 (x) x0	y1=105.724+ 1.373 (x)x1	y2= 106.64 + 1.35 (x) x2	y0= 103.76+ 1.43 (x) x0
Predicted ht (y)	164.484cm	164.478cm	164.478cm	155.289cm	155.289cm	155.289cm

\*\*\* Significant at  $p<0.0001$ ; rt= right; lt= left.

We have also estimated the multiplication factor (M.F.) for PCTL (Table 4). The average M.F. was found to be 4.302 in male and 4.306 in female. With the help of this multiplication factor the average stature was calculated as 164.5cm for male and 155.292cm for female, which showed the average error of 0.00 cm in male and 0.008 cm in female.

**Table 4:** Multiplication factor (M.F.) in both gender for tibial length

Tibia	Male		Female	
	Rt	Lt	Rt	Lt
PCTL	38.26 cm	38.22 cm	36.099 cm	36.03 cm
M.F.	4.299	4.304	4.302	4.3099
Average M.F.	4.302		4.306	
Calculated average Stature	164.5cm		155.292cm	

The stature estimated from PCTL with the help of formulated M.F was compared with stature estimated by

regression formula, the average error was found to be 0.022cm in male and 0.003cm in females. The average error was nearly insignificant and less than 1cm; hence a multiplication factor can also be used as a formula for estimation of stature.

## DISCUSSION

M.F or regression formulae are regional population and gender specific and thus provide greater reliability in estimated stature.<sup>8,19</sup> We have also derived regression formula and M.F both. Kaore et al<sup>16</sup> reported that the Regression formulae are more dependable than multiplication factor for estimation of stature. Kate and Muzumdar<sup>28</sup> after comparing the derived regression equation for Maharashtrian and Punjabis with that of Pearson's regression formula derived from English bone stated that Pearson's regression equation does not give exact results in Indian population. Similar view by Kaore et al.<sup>16</sup> They suggested that the regression formula derived by Allbrook<sup>7</sup> for estimating the stature in the British population is not suitable to estimate the stature in Indian population.<sup>16,28</sup>

The union of the epiphysis and diaphysis commonly occurs at the age of 18-20 years and after that individuals stop growing in height, therefore all the individuals considered for the purpose of the study were either at or above the age of 20 years. In this study, the mean height for male was 164.5±8.257cm and for female was 155.3±5.854cm; and the mean PCTL for male was 38.24±2.343cm which was significantly ( $p < 0.0001$ ) greater than the female which was 36.064±2.464cm. Results of the present study are similar to the findings of Yayim Yili,<sup>29</sup> Agnihotri et

al,<sup>30</sup> Chavan et al,<sup>23</sup> Bhavna and Surinder Nath<sup>8,19</sup> and many others, who observed that there was no statistically significant difference in the length of right and left tibia in both males and females. Mukta Rani<sup>31</sup> compared the bilateral percutaneous measurement of tibia and expressed that left tibia is longer than the right tibia in both sexes. Allbrook<sup>7</sup> in 1961, compared both estimated stature derived from length of dried tibia and from the average percutaneous tibial length. There was no difference in stature estimated from two different sets of tibia. The average stature was 170.06 cm for British male population. Chavan et al<sup>23</sup> estimated the mean height of male and female to be 167.89 cm± 6.21 cm and 151.41 cm± 5.04 cm respectively. Mean PCTL was 37.32cm ±2.18 cm for male and 34.44cm ± 2.10 cm for female. Mukta Rani et al<sup>31</sup> estimated the stature in students of Delhi to be 169.5cm in male and 159.5cm in females, which were higher than our results. Kaore et al<sup>16</sup> estimated average stature 170.089cm for Indian male population with an average error less than 1cm. Bhavna and Surender Nath<sup>8</sup> in their study on male Shia Muslims in Delhi derived the following linear regression equation; Height in cms =  $84.74 + 2.27x$  (PCTL) ± 3.67, which is comparable to our study, but exemplifies the fact that the regression equation derived will be population group/region specific<sup>15</sup>. In our study, we assessed both males and females, which has not been done in the above study. Our estimated stature nearly correlates well with that of Bhavna and S. Nath<sup>8,19</sup> who estimated stature to be 167.66 cm for males and 154.40 cms for females.

According to Trotter and Gleser<sup>32</sup> world population is getting taller and therefore

the relationship between height and length of long bones is changed and fresh formulae or M.F are needed for each generation, hence they attempted to find out fresh M.F for Indians. Our values of multiplication factor are comparable with those of Bhavana and Surinder Nath<sup>8,19</sup> who gave the values for M.F as 4.60 in males and 4.59 in females. The M.F in our study was 4.302 for male and 4.306 for female. Chavan et al<sup>23</sup> estimated the average M.F for tibia to be 4.77 in male and 4.88 in female and the average stature calculated 170.69 cm for male and 157.06 for females, which showed the average error of 0.61 cm in male and 0.86 cm in females. Chavan et al<sup>23</sup> estimated the value of 'r' for males was 0.82 and for females 0.68. Both these values were statistically significant. Bhavana and Surinder Nath<sup>8</sup> estimated  $r=0.765$  for male. In our study  $r=0.4342$  for males, which is smaller than other studies and  $r=0.6014$  for females, which nearly correlates with other studies. Petrovečki et al<sup>4</sup> tested a new radiographic approach to the stature prediction that could be used in the identification process of human skeletal remains of unknown identity.

The stature of 19 female and 21 male adult cadavers was measured within 24 hours after death and considered equal to the living stature. The anteroposterior radiographs of all limbs were taken and the maximum length of the six long bones was measured from radiographs. There was a significant difference in the stature and maximum length of long bones between female and male cadavers ( $p<0.001$  for all). The correlation between the stature and long bone length was best for the humerus in females ( $r=0.792$ ) and the tibia in males ( $r=0.891$ ). Regression equations

specific to Croatian population were computed separately for each long bone in males and females and proven reliable in predicting the living stature of the individual.

## CONCLUSION

There was no significant difference in the per-cutaneous length of right and left tibia in both genders, thus showing bilateral symmetry in the length of Tibia in both genders. In both genders stature estimated by regression formulae for per-cutaneous tibial length of people of Gwalior region was similar to average measured stature with an error of less than 1cm which was statistically insignificant  $P > 0.05$ . Multiplication factor for length of tibia was similar to average measured stature with an error of less than 1cm. This was statistically insignificant  $P > 0.05$ . It was concluded that it is possible to determine the stature of a deceased person whose only body part available is a mutilated leg, by using the data and formula derived from the present study fairly accurately to some extent. However the formulae derived cannot be generalized to all population groups, hence it is necessary to derive regression equations which are region wise and population specific. Thus the data of this study are recommended in anthropological studies for stature estimation amongst the ethnic group under study. Further scope of this study can be expended by including the subjects who are not native of the Ajmer region.

**Source of Funding:** - Self

**Interest of conflict:** - None.

## References

1. Trotter M, Gleser GC. Estimation of stature from long bones of American

- whites and negroes. *Am J Phys Anthropol* 1952;10:463-514.
2. Mall G, Hubig M, Büttner A, Kuznik J, Penning R, Graw M. Gender determination and estimation of stature from the long bones of the arm. *Forensic Sci Int* 2001;117: 23-30.
  3. Duyar I, Pelin C. Body height estimation based on tibia length in different stature groups. *Am J Phys Anthropol* 2003;122:23-7.
  4. Petrovečki V, Mayer D, Šlaus M, Strinović D, Škavić J. Prediction of stature based on radiographic measurements of cadaver long bones: a study of the Croatian population. *Journal of Forensic Sciences*. 2007; 52 (3):547-52.
  5. Lundy JK, Feldesman MR. Revised equations for estimating living stature from long bones of the South African Negro. *S Afr J Sci* 1987; 83: 54-55.
  6. Brothwell D, Zakrzewski. Metric and non-metric studies of archaeological human bone. In: Brickley M, McKinley JI, editors. *Stature estimation. Guidelines to the Standards for recording Human Remains*. Southamp-ton/Reading: BABAO/Institute of Field Archeologists; 2004.
  7. Allbrook D. The estimation of stature in British and East African males based on tibial and ulnar bone lengths. *J. For. Med.*1961; 8: 15-28.
  8. Bhavna S, Surinder Nath. Estimation of Stature on the Basis of Measurements of the Lower Limb. *Anthropologist Special Volume*.2007; 3: 219-222.
  9. Ozaslan A, Iscan MY, Ozaslan I, Tugcu H, Koc S. Estimation of stature from body parts. *Forensic Sci Int* 2003;132:40-5.
  10. Badkur P, Nath S. Use of regression analysis in reconstruction of maximum bone length and living stature from fragmentary measures of the ulna. *Forensic Sci Int* 1990;45:15-25.
  11. Holland TD. Estimation of adult stature from fragmentary tibias. *J Forensic Sci* 1992;37:1223-9.
  12. Prasad R, Vettivel S, Jeyaseelan L, Isaac B, Chandi G. Reconstruction of femur length from markers of its proximal end. *Clin Anat* 1996;9:28-33.
  13. Trotter M, Gleser GC. Estimation of stature from long bones of American Whites and Negroes *Am. J. Phys. Anthropology*. 1952; 10:463-514.
  14. Trotter M, Gleser GC. A re-evaluation of estimation of stature based on measurements of stature taken during life and of long bones after death. *Am. J Phys. Anthropology*.1958; 16:79-123.
  15. Lal CS, Lala JK. Estimation of height from tibial and ulnar lengths in North Bihar. *Journal of Indian Medical Association* .1972; 58(4):120-121.
  16. Kaore A, Kaore BP, Kamdi A , Kaore S. Stature Estimation From Tibial Length. *NJIRM*. 2012; 3(2):51-56.
  17. Krici Y, Ozan H. Determination of gender from the tibia of adult Turkish cadavers. *Kaibogaku Zasshi*. 1999; 45:537-543.
  18. Cox HWV. *Medical jurisprudence and Toxicology*. 6<sup>th</sup> ed. The law book company.1990;171-174.
  19. Bhavna S, Surinder Nath. Use of Lower Limb Measurements in Reconstructing Stature among Shia Muslims. *The internet Journal of Biological Anthropology*. 2009; 2(2).
  20. Steele DG. Estimation of Stature from Fragments of Long limb bones, in: Stewart. T.D (ed.) *Personal Identification in Mass Disasters*.

- Smithsonian Institute: Washington D.C, 1970 ; 85-97.
21. Nath BS. Estimation of stature from long bones in Indians of United Provinces: A Medico-Legal enquiry in Anthropometry. *Ind. J. Med Res* 1931; 18: 1245-1253.
  22. Kumar P, Shahnawaz K, Varma G. Study of Estimation of Stature by the Length of Femur. *Journal of Evolution of Medical and Dental Sciences* 2014; 3(12): 3166-72.
  23. Chavan SK, Chavan KD, Mumbre SS, Makhani CS. Stature and Percutaneous Tibial Length: A Correlational Study in Maharashtrian Population. *Indian Journal of Forensic Medicine and Pathology*. 2009; 2(3):109-12.
  24. Siddiqui MAH, Shah MA. Estimation of Stature from long bones of Punjabis. *Ind. J. Med. Res.* 1944; 32: 105-108.
  25. Iscan MY. Global forensic anthropology in the 21st century (Editorial). *Forensic Sci Int*, 2001 117: 16.
  26. Iscan MY. Forensic anthropology of gender and body size (Editorial). *Forensic Sci Int*, 2005 147: 107-112.
  27. Lal CS, Lala JK. Estimation of height from tibial and ulnar lengths in North Bihar. *Journal of Indian Medical Association*.1972; 58(4):120-121.
  28. Kate BR, Majumdar RD. Stature estimation from femur and humerus by regression and autometry. *Acta. Anat.* 1976; 94: 311-320. [29] Yayim Yili. Estimation of stature from tibial length. *Journal of Forensic Medicine*. 1996; 12: 87-93.
  29. Agnihotri AK, Kachhwaha S, Jowaheer V, Singh AP. Estimating stature form percutaneous length of tibia and ulna in indo-Mauritian population. *Forensic Science International*. 2009;187 (1) : 109.e1-109.e3.
  30. Rani Mukta , Tyagi AK, Verma SK, Kohli A. Estimation of stature from percutaneous measurements of legs (1999-2000). *Journal of Forensic Medicine and Toxicology* 2004; 21(1):12-14.
  31. Trotter M, Gleser GC. Estimation of stature from long bones of American Whites and Negroes. *Am. J. Phys. Anthropology*. 1952; 10:463-514.